# Published for OXFORD INTERNATIONAL AQA EXAMINATIONS

# International A Level FURTHER MATHEMATICS with Statistics



Brian Gaulter Mark Gaulter Brian Jefferson John Rayneau



# Contents

## **AS Pure**

## 1 Loci, Graphs and Algebra

1.1	Loci	2
1.2	Rational functions	3
1.3	Conic sections	11
Sum	imary	16
Revi	ew exercises	17
Prac	tice examination questions	17

## 2 Complex Numbers

2.1	What is a complex number?	20
2.2	Calculating with complex numbers	22
2.3	Argand diagram	25
2.4	Loci in the complex plane	28
Sum	mary	31
Revi	ew exercises	32
Prac	tice examination questions	33

## **3** Roots and Coefficients of a Quadratic Equation

	-	
3.1	Roots of a quadratic equation	
3.2	Finding an equation with roots that are	$(\mathbb{Z})$
	a function of existing roots	
Sum	mary	39
Revi	ew exercises	39
Prac	tice examination questions	

#### 4 Series

4.1 Su	mmation formulae [		
4.2 Me	thod of differences .		45
Summa	ry	<u> </u>	
Review	exercises	<u>,                                     </u>	47
Practice	examination question	ons	

## 5 Trigonometry

General solutions of	
trigonometric equations	49
Solving equations involving more	
complicated terms	53
mary	55
ew exercises	56
Practice examination questions5	
	General solutions of trigonometric equations Solving equations involving more complicated terms mary ew exercises tice examination questions

## 6 Calculus

6.1	Gradient of a tangent to a curve	58
6.2	Rates of change	60

6.3 In	mproper integrals	63
Summ	ary	65
Review	v exercises	66
Practic	e examination questions	66

## 7 Matrices and Transformations

7.1	Introduction to matrices	68
7.2	Transformations	72
7.3	Common transformations	76
7.4	Invariant points and lines	78
Sum	mary	80
Revie	ew exercises	81
Prace	tice examination questions	82

## 8 Linear graphs

8.1 Relationship between data	84
<b>8.2</b> When the power of <i>x</i> is unknown, or when	
x is in the exponent	87
Summary	90
Review exercises	91
Practice examination questions	92

## 9 Numerical Methods

9.1	Solutions of polynomial equations	
9.2	Step-by-step solution of differential	
	equations	
Sum	mary	101
Revi	ew exercises	
Prac	tice examination questions	

## **AS Statistics**

## 10 Bayes' Theorem

10.1 Tree diagrams	104
10.2 Bayes' Theorem	109
Summary	112
Review exercises	112
Practice examination questions	113

#### 11 Discrete Uniform and Geometric Distributions

<b>11.1</b> Discrete uniform distribution	116
<b>11.2</b> Geometric distribution	121
Summary	125
Review exercises	126
Practice examination questions	127

## 12 Probability Generating Functions A2 Pure

12.1	Probability generating functions and their	
	properties	131
12.2	Some standard distributions	135
12.3	Sums of independent random	
	variables	139
Sum	mary	144
Revi	ew exercises	144
Prac	tice examination questions	145

#### **13 Linear Combinations of Discrete Random Variables**

13.1	Linear combinations of discrete	
	random variables	148
13.2	Independent discrete random	
	variables	153
Sum	mary	155
Revie	ew exercises	155
Prace	tice examination questions	156

## **AS Mechanics**

#### 14 Constant Velocity in Two **Dimensions**

14.1 Vectors in component form	158
<b>14.2</b> Position, displacement, distance, velocity	
and speed	159
14.3 Resultant velocity	162
14.4 Relative velocity, closest approach	
and interception	165
Summary	170
Review exercises	171
Practice examination questions	171

## 15 Dimensional Analysis

<b>15.1</b> Dimensions and dimensional	
consistency	174
15.2 Finding a formula	177
Summary	179
Review exercises	180
Practice examination questions	181

Y

#### **16 Collision in One Dimension**

16.1 Impulse and momentum	182
<b>16.2</b> The principle of conservation of linear	
momentum	186
16.3 Elastic impact	188
16.4 Multiple collisions	193
Summary	195
Review exercises	196
Practice examination questions	196

#### **17 Roots and Polynomials**

17.1 Roots of higher-order equations	198
17.2 Complex roots of a polynomial	
equation	201
Summary	
Review exercises	205
Practice examination questions	206

#### **18 Proof by Induction and Finite** Series

<b>18.1</b> Proof by induction		)8
18.2 Finite series	21	15
Summary	21	17
Review exercises	21	17
Practice examination quest	ions 21	17

## 19 Series and Limits

19.1 Convergence	220
19.2 Maclaurin's theorem	221
19.3 Finding the limits of a series	227
19.4 Improper integrals	228
Summary	230
Review exercises	231
Practice examination questions	231

## 20 De Moivre's Theorem

20.1	De Moivre's theorem	234
20.2	Further applications of de Moivre's	
	theorem to complex numbers	239
20.3	Trigonometric identities	245
Sumi	mary	252
Revie	ew exercises	253
Pract	ice examination questions	253

## **21 Polar Coordinates**

<b>21.1</b> Position of a point	256
<b>21.2</b> Sketching curves given in polar	
coordinates	259
<b>21.3</b> Area of a sector of a curve	262
Summary	266
Review exercises	267
Practice examination questions	267

#### 22 The Calculus of Inverse **Trigonometric Functions**

22.1 Inverse trigonometric functions	270
22.2 Differentiation and integration of inv	verse
trigonometric functions	
Summary	279
Review exercises	280
Practice examination questions	

#### 23 Arc Length and Area of Surface of Revolution

23.1 Arc length	282
<b>23.2</b> Area of a surface of revolution	285
Summary	286
Review exercises	287
Practice examination questions	287

#### 24 Hyperbolic Functions

24.1 Hyperbolic functions	288
24.2 Differentiation of hyperbolic functions	292
24.3 Inverse hyperbolic functions	294
Summary	303
Review exercises	304
Practice examination questions	305

#### 25 Differential Equations of First and Second Order

25.1	First-order linear equations	306
25.2	Second-order differential equations	308
25.3	Using a complementary function and particular	r
	integral to solve a first order equation	318
Sum	mary	318
Revie	ew exercises	319
Pract	tice examination questions	319

#### 26 Vectors and Three-Dimensional Coordinate Geometry

26.1 Vectors	320
26.2 Vector product	324
<b>26.3</b> Applications of vectors to	
coordinate geometry	329
26.4 Equation of a line	329
26.5 Equation of a plane	329
26.6 Scalar triple product and its applications	338
Summary	342
Review exercises	344
Practice examination questions	344

## **27 Solutions of Linear Equations**

27.1	Simultaneous linear equations	346
27.2	Using determinants to find the number of	
	solutions of three simultaneous equations	349
Sum	mary	351
Revie	ew exercises	352
Pract	tice examination questions	352

## 28 Matrix Algebra

28.1 Inverse matrices	354
28.2 Transformations	356
28.3 Invariant points and lines	361
Summary	367
Review exercises	368
Practice examination questions	368

## **A2 Statistics**

#### **29 Moment Generating Functions**

29.1	Moment generating functions of	
	random variables	370
29.2	Moment generating functions of some	
	standard distributions	375
<b>29.3</b>	Sums of independent random variables	379
Sum	mary	381
Revie	ew exercises	382
Prace	tice examination questions	383

## 30 Estimators

30.1 Review of key concepts	384
<b>30.2</b> Estimators and estimates	386
Summary	393
Review exercises	393
Practice examination questions	394

## 31 Estimation

<b>31.1</b> Confidence intervals for the mean	
of a distribution	398
<b>31.2</b> Finding the sample size for a given	
confidence interval width	407
31.3 Inferences from confidence intervals	409
Summary	412
Review exercises	412
Practice examination questions	413

#### 32 Further Hypothesis Testing 1 – Means and Variances

32.1	Power of a hypothesis test	415
32.2	Tests for the difference between the mean	
	of two independent distributions	419
32.3	Tests for the difference between the means	
	of two non-independent distributions	428
<b>32.4</b>	Tests related to population variances	431
Sum	mary	438
Revie	ew exercises	439
Pract	tice examination questions	440

#### 33 Further Hypothesis Testing 2 – Goodness of Fit Tests and Tests of No Association

<b>33.1</b> Goodness of fit tests	442
<b>33.2</b> Tests of no association	450
Summary	455
Review exercises	455
Practice examination questions	456
Answers	460
Index	497

# Loci, Graphs and Algebra

#### Introduction

Polynomial functions always form a continuous curve with no breaks. However, when you divide one polynomial by another, the graph of the new function can have breaks in it and is said to be **discontinuous**. An example of such graphs is a **conic section**, which is the curve formed when a plane intersects a right circular cone. Some of the curves you meet the most in the real world are examples of conic sections, such as the ellipse that describes Earth's orbit around the Sun and the parabola that models the path of a football.

#### Recap

You will need to remember how to ...

- ► Solve construction problems involving loci.
- ► Solve equations, including quadratics.
- Sketch basic graphs such as  $y = x^2$ .
- ► Find the distance between two points in Cartesian coordinates.
- ► Solve simple inequalities such as 4x + 7 > 3(x 4) and  $x^2 7x + 10 \ge 0$ .
- ► Transform graphs using stretches, reflections and translations.

## 1.1 Loci

In the context of graphs, a **locus** (plural **loci**) is a set of points that follow a given rule. Therefore, a locus can be represented by an equation.

For example, a locus is given as the points that are a distance of four units from the point (2, 3). This locus forms a circle with centre (2, 3). You know from previous studies that the equation of a circle is given in the form  $(x-a)^2 + (y-b)^2 = r^2$ , where *r* is the radius of the circle with centre (*a*, *b*). Therefore, the locus described above can be given as the Cartesian equation  $(x-2)^2 + (y-3)^2 = 16$ .

To find the Cartesian equation of a given locus, consider a general point on the curve (x, y) and use what you know about loci to help you formulate an appropriate equation.

#### **Example 1**

Find the Cartesian equation of the locus of points that are equidistant from the point (-1, 4) and the line x = 2.

#### **Objectives**

By the end of this chapter, you should know how to:

- Sketch graphs of rational functions.
- Find equations of asymptotes to graphs.
   Solve inequalities
- involving rational functions.

Describe and sketch various conic sections.

- Find the points of intersection between conic sections and coordinate axes and various straight lines.
- Find the Cartesian equation of simple loci that are described verbally.



(x, y) is a general point that obeys the rule. Distance of (x, y) from line x = 2 is |x - 2|.

Distance from (x, y) to (-1, 4) is

 $\sqrt{(x+1)^2 + (y-4)^2}$  $(x+1)^2 + (y-4)^2 = (x-2)^2$ 

 $x^{2}+2x+1+y^{2}-8y+16=x^{2}-4x+4$ Equation of the locus is  $(y-4)^{2}=-6x+3$ .

#### **Exercise 1**

- Find the Cartesian equation of the locus of points which are equidistant from the point (3, -2) and the line y = 5.
- Find the Cartesian equation of the locus of points which are a distance of four units from the point (2, -3).
- Find the Cartesian equation of the locus of points which are equidistant from the point (-5, 3) and the line x = 2.
- Find the Cartesian equation of the locus of points which are a distance of 4√2 units from the point (4, −4).

## **1.2 Rational functions**

During your A-level Mathematics studies you will have learned how to sketch curves. In this chapter, you will learn how to sketch curves for functions that are more complicated than trigonometric or polynomial ones.

The graph of a rational function will always have a horizontal asymptote provided that the degree of *x* in the denominator is the same as or larger than the degree of *x* in the numerator. In this chapter, you will only deal with cases where there is a horizontal asymptote.

# Sketching rational functions with a linear denominator

In order to sketch rational functions equations of the form  $y = \frac{ax+b}{cx+d}$ , you need to start by finding the asymptotes.

You should remember that in the context of sketching a curve, an **asymptote** is a line that becomes a tangent to the curve as *x* or *y* tends to infinity. (Vertical asymptotes are the lines where the graph is undefined.)

For example, take the curve of  $y = \frac{4x - 8}{x + 3}$ .

In order for  $y \rightarrow \pm \infty$ , the denominator of this function must tend to zero.

That is, as  $x + 3 \rightarrow 0$ ,  $x \rightarrow -3$ . Hence, x = -3 is an asymptote.

To find the asymptote as  $x \rightarrow \pm \infty$ , you express the function as

$$y = \frac{4 - \frac{6}{x}}{1 + \frac{3}{x}}$$
  
As  $x \to \pm \infty$ ,  $\frac{3}{x} \to 0$  and  $\frac{8}{x} \to 0$ .  
Therefore,  $y \to \frac{4}{1} = 4$ .

#### Note

The two distances are equal.

#### Note

You will discover later in this chapter that this is a **conic equation** of a parabola.

Tip

The asymptote occurs where the graph is undefined, so to find the vertical asymptote you need to equate the denominator of the rational function to zero.

#### Tip

Divide the top and bottom by *x*.

Hence, y = 4 is also an asymptote.

Notice that as  $x \to \pm \infty$ , the largest terms in the numerator and the denominator are 4x and x respectively, and so  $y \approx 4x \div x = 4$ .

x = -3 is a **vertical asymptote**, as it is parallel to the *y*-axis.

*y* = 4 is a **horizontal asymptote**, as it is parallel to the *x*-axis.

To be able to sketch  $y = \frac{4x-8}{x+3}$ , you also need to find where it crosses the *x*- and *y*-axes.

When x = 0:  $y = -\frac{8}{3}$  When y = 0:  $4x - 8 = 0 \implies x = 2$ In summary, you proceed as follows:

- **1.** Draw the asymptotes using dashed lines.
- **2.** Mark the points where the curve crosses the axes; as the numerator and the denominator of the function each contain only a linear term in *x*, the curve cannot cross either asymptote.
- **3.** Considering the curve for x > -3, you can see that *y* tends to  $-\infty$  as *x* approaches -3 from values of *x* greater than -3. Hence, the curve tends to  $+\infty$  as *x* approaches -3 from values of *x* less than -3.
- 4. You can now complete the curve of  $y = \frac{4x-8}{x+3}$ .

#### To sketch the curve of a rational function in the form y = 4

- 1. Find the vertical asymptote by equating the denominator to zero; find the horizontal asymptote by dividing the numerator and denominator by *x*, that is, expressing the function as  $y = \frac{a + \frac{b}{x}}{c + \frac{d}{x}}$  and then considering what happens when *x* tends to infinity.
- 2. Substitute x = 0 and y = 0 into the function to find where the curve crosses the axes.
- 3. If necessary, consider the curve for the x-values, to see what happens to the y-values as x tends to  $\pm \infty$ .

#### Example 2



Vertical asymptote: as  $x \to \pm \infty$ ,  $y \to \frac{2}{1}$ , y=2Vertical asymptote: as  $y \to \pm \infty$ ,  $x-5 \to 0$ , x=5When x=0:  $y = \frac{-6}{-5} = \frac{6}{5}$ 





When the degrees of the numerator and denominator are the same, you can find the horizontal asymptote by dividing the leading coefficient of the numerator by the leading coefficient of the denominator.

n

2

 $\hat{x}$ 

x



-3

First, find the asymptotes.

#### Note

x

5

Next, find where the curve crosses the axes.



When  $y = 0: 2x - 6 = 0 \implies x = 3$ 

Note that the asymptotes here are parallel to the coordinate axes.

#### Exercise 2

In questions 1, 2 and 3, state the equations of the asymptotes for the curve.

- 2  $y = \frac{5}{x+2}$ 1  $y = \frac{x-1}{2x+6}$ **3**  $y = \frac{2x+7}{x-3}$ 4 Show that y = 2 is an asymptote for the curve  $y = \frac{2x+3}{x+4}$ . Show that y = -4 is an asymptote for the curve  $y = \frac{8-4x}{x+2}$ .
- 6 Sketch  $y = \frac{6x-3}{x+4}$ . **7** Sketch  $y = \frac{3x-6}{x-1}$ . 8 Sketch  $y = \frac{2x+8}{3x-5}$ , stating the equations of the asymptotes of the curve.

## Sketching rational functions with a quadratic denominator

When sketching rational functions with equations of the form  $y = \frac{ax^2 + bx + c}{dx^2 + ex + f}$ where both the numerator and the denominator are quadratics, you proceed as before by finding the asymptotes, where the graphs cross the axes, and consider the shape of the curve. However, with these functions, the shape of the curve requires consideration of stationary points and the number of asymptotes depends on the number of roots of the quadratic denominator.

- Two different roots will result in two vertical asymptotes.
- One repeated real root will result in one vertical asymptote.
- No roots indicates that there are no vertical asymptotes.

Regardless of the number of asymptotes, there may be values of y that are not realised by any value of x. Therefore, you need to find the **range** of the rational function in order to determine the set of possible values of y and in turn find the maximum and minimum points (the stationary points) of the curve.

## Example 3

Find the range of possible values of y when  $y = \frac{3x-4}{x^2+3x-4}$ .

```
Cross-multiplying,
```

```
yx^{2} + 3yx - 4y = 3x - 4

\Rightarrow yx^{2} + (3y - 3)x + 4 - 4y = 0
```

## Therefore,

 $(3y-3)^2 - 4y(4-4y) \ge 0$ 

#### Note

To find the range of values of  $\gamma$ , you need to find the values for which  $y = \frac{3x-4}{x^2+3x-4}$  has real solutions for x.

## Note

Now copy and complete the sketch of  $y = \frac{2x-6}{x-5}$  by also considering what will happen to the *v*-values as the x-values change.

(continued)

5

For x to be real, we know that

Note

 $b^2 - 4ac \ge 0$ .

In questions **5** to **7**, sketch the curve and clearly mark the points where the curve crosses the coordinate axes.

5 
$$\frac{x^2}{36} + \frac{y^2}{25} = 1$$
  
6  $\frac{(x-4)^2}{25} + \frac{(y-3)^2}{16} = 1$   
7  $\frac{(x-2)^2}{25} - \frac{(y+6)^2}{16} = 1$ 

8 A straight line through (1, 0) with gradient *m* intersects the hyperbola  $\frac{x^2}{9} - \frac{y^2}{25} = 1$  at point *P*. Show that the *x*-coordinate of point *P* satisfies the equation  $(25 - 9m^2)x^2 + 18m^2x - (9m^2 + 225) = 0$ .

- 9 Write the asymptotes of (x 5)(y + 3) = 6, and sketch the curve.
- 10 An ellipse has the equation  $\frac{x^2}{4} + \frac{y^2}{25} = 1$ .
  - **a** Sketch the ellipse.
  - **b** Given that the line y = x + k intersects the ellipse at two distinct points, show that  $-\sqrt{29} < k < \sqrt{29}$ .
  - **c** The ellipse is translated by the vector  $\begin{pmatrix} a \\ b \end{pmatrix}$  to form another ellipse whose equation is

$$25x^2 + 4y^2 + 50x - 24y = c$$

Find the values of *a*, *b* and *c*.

## Summary

- ► A locus is a set of points that obey a certain rule, and a locus can be expressed graphically, verbally or in the form of an equation.
- Asymptotes show the 'end behaviour' of a graph as x or  $y \to \pm \infty$ .
- To sketch the graphs of rational, parabola, ellipse and hyperbola equations you might need to find the:
  - Asymptotes (if applicable; parabolas and ellipses do not have asymptotes)
  - Intercepts with the axes, if any
  - Coordinates of any maxima or minima (if applicable).
- ► You can solve a rational inequality by:
  - Using algebra to multiply both sides of the inequality by  $(cx + d)^2$
  - Sketching  $y = \frac{ax+b}{cx+d}$ , then solving  $\frac{ax+b}{cx+d} = k$  and comparing the two results to find the solution.
- ► Conic sections are a family of curves with standard equations, and include the:
  - Parabola,  $y^2 = 4ax$
  - Ellipse,  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$
  - Hyperbola,  $\frac{x^2}{a^2} \frac{y^2}{b^2} = 1$
  - Rectangular hyperbola,  $xy = c^2$ .

► The transformations of more complicated curves follow the same rules linear transformations do.

If y = f(x), then

- y = f(x) + a results in a positive translation in the *y*-direction
- y = f(x) a results in a negative translation in the *y*-direction
- y = f(x + a) results in a negative translation in the *x*-direction
- y = f(x a) results in a positive translation in the *x*-direction
- y = af(x) results in a stretch parallel to the *y*-axis, with scale factor *a*
- y = f(ax) is a stretch parallel to the x-axis, with scale factor  $\frac{1}{a}$
- y = -f(x) is a reflection in the *y*-axis
- y = f(-x) is a reflection in the *x*-axis.

#### **Review exercises**

- Find the Cartesian equation of the locus of points which are equidistant from the point (5, −1) and the *y*-axis.
- 2 Sketch the graph of  $y = \frac{2x^2 + 3x 5}{x^2 x 2}$
- 3 Sketch the graph of  $y = \frac{3x^2 + 4x + 4}{x^2 2x 3}$

4 Solve 
$$\frac{2x-1}{x+3} > 3$$
.

5 Solve 
$$\frac{x^2 - x - 2}{x^2 + 3x + 2} > 1.$$

6 Sketch 
$$\frac{(x-3)^2}{36} - \frac{(y+2)^2}{25} = 1$$
.

7 The curve  $x^2 + \frac{y^2}{9} = 1$  is translated by k units in the positive y-direction.

- **a** Show that the equation of the curve after this translation is  $x^2 + \frac{(y-k)^2}{2} = 1.$
- **b** Show that if the line x + y = 3 intersects the translated curve, the *y*-coordinate of the points of intersection satisfies the equation  $10y^2 (54 + 2k)y + k^2 + 72 = 0$ .

## **Practice examination questions**

**a** i Write down the equations of the two asymptotes of the curve y = 1/(x-3). (2 marks)
 ii Sketch the curve y = 1/(x-3), showing the coordinates of any points of intersection with the coordinate axes. (2 marks)

- iii On the same axes, again showing the coordinates of any points of intersection with the coordinate axes, sketch the line y = 2x 5. (1 mark)
- **b** i Solve the equation  $\frac{1}{x-3} = 2x-5$ . (3 marks) ii Find the solution of the inequality  $\frac{1}{x-3} < 2x-5$ . (2 marks)

AQA MFP1 June 2010

17



18

iii Hence show that the curve has only one stationary point and find its coordinates. (No credit will be given for solutions based on differentiation.) (4 marks) c Sketch the curve and its asymptotes. (3 marks) AQA MFP1 June 2013 5 An ellipse is shown on the right. The ellipse intersects the *x*-axis at the points *A* and *B*. The equation of the ellipse is  $\frac{(x-4)^2}{4} + y^2 = 1$ . **a** Find the coordinates of *A* and *B*. (2 marks) **b** The line y = mx (m > 0) is a tangent to the ellipse, with point of contact Р. C  $\hat{x}$ i Show that the *x*-coordinate of *P* satisfies the equation  $(1+4m^2)x^2-8x+12=0$ (3 marks) **ii** Hence find the exact value of *m*. (4 marks) iii Find the coordinates of P. (4 marks) AQA MFP1 January 2013 **6** The curve *C* has equation  $y = \frac{x}{(x+1)(x-2)}$ . The line *L* has equation  $y = -\frac{1}{2}$ . **a** Write down the equations of the asymptotes of *C*. (3 marks) **b** The line *L* intercepts *C* at two points. Find the *x*-coordinates of these two points. (2 marks) **c** Sketch *C* and *L* on the same axes. (You are given that the curve C has no stationary points). (3 marks) **d** Solve the inequality (3 marks) AQA MFP1 June 2012